

Microcausality and Energy-Positivity in all frames imply Lorentz Invariance of dispersion laws

by

Jacques Bros¹ and Henri Epstein²

¹ Service de Physique Théorique, CEA-Saclay, F-91191 Gif-sur-Yvette, France

² Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France

Abstract. A new presentation of the Borchers-Buchholz result of the Lorentz-invariance of the energy-momentum spectrum in theories with broken Lorentz symmetry is given in terms of properties of the Green's functions of microcausal Bose and Fermi-fields. Strong constraints based on complex geometry phenomenons are shown to result from the interplay of the basic principles of causality and stability in Quantum Field Theory: if microcausality and energy-positivity in all Lorentz frames are satisfied, then it is unavoidable that all stable particles of the theory be governed by Lorentz-invariant dispersion laws; in all the field sectors, discrete parts outside the continuum as well as the thresholds of the continuous parts of the energy-momentum spectrum, with possible holes inside it, are necessarily represented by mass-shell hyperboloids (or the light-cone). No violation of this geometrical fact can be produced by spontaneous breaking of the Lorentz symmetry, even if the field-theoretical framework is enlarged so as to include short-distance singularities wilder than distributions.

1- Introduction

In a recent work [1], it has been advocated that the occurrence of spontaneous Lorentz and CPT violations in Quantum Field Theories governed by suitable *non-local* Lagrangians can very well generate *non-Lorentz-invariant dispersion laws* ⁽¹⁾ which avoid the problems with stability and causality. Such Lorentz violation effects produced at Planck scale might then in principle be observed at lower energies in particle physics. In support of their claim, the authors of [1] have produced examples of possible “non-local models” in which the quadratic part of the Lagrangian corresponds to a dispersion law $p_0 = \omega(\vec{p})$ enjoying the following properties:

a) The hypersurface $\mathcal{M}$ with equation $p_0 = \omega(\vec{p})$ differs from a Lorentz-invariant mass shell hyperboloid,

b) $\mathcal{M}$ is contained in the positive energy-momentum cone $\overline{V}^+$ ($p_0 \geq |\vec{p}|$),

c) For every momentum $\vec{p}$, the “group velocity condition” $|\partial\omega(\vec{p})| \leq 1$ holds, which means that $\mathcal{M}$ admits a space-like (or light-like) tangent hyperplane at each of its points.

⁽¹⁾ We prefer keeping here the terminology of “dispersion law” (used traditionally e.g. in Thermal Quantum Field Theory) rather than adopting the new usage of “dispersion relation”, which is of course confusing in a domain where (Cauchy-type) dispersion relations relating the absorptive and dispersive parts of Feynman-type amplitudes remain a basic tool of frequent use.

While condition b) expresses energy-positivity in all Lorentz frames, condition c) ensures that all wave-packets satisfying the dispersion law $p_0 = \omega(\vec{p})$ propagate “essentially” with a *subluminal (or luminal) velocity*; essentially means “up to the quantum spreading of wave-packets, of the order of the Planck constant”, as it is the case for the solutions of the Klein-Gordon and Dirac equations. However, in spite of this causal interpretation in terms of particle wave-packets, condition c) should by no means be taken as a criterion of *microcausality* for the underlying Quantum Field Theory. Microcausality states that the commutator (resp. anticommutator) $[\Phi(x), \Phi(x')]_{\mp}$ of a boson (resp. fermion) field $\Phi(x)$ should vanish in the whole region of *relativistic spacelike separation* $\{(x, x'); (x - x')^2 < 0\}$. As we shall see below, the “group velocity condition” c) only appears as a necessary consequence of microcausality, but the converse is not true. This is why, without underestimating the interest of the mentioned examples of [1] satisfying conditions a), b) and c), it appeared worthwhile to us to warn the field-theorist community that such examples cannot correspond to field models satisfying microcausality. In fact, at a high level of generality including field theories with local singularities of arbitrary strength (i.e. wilder than distributions) in spacetime, the requirement of microcausality represents such a strong constraint that, when combined with energy positivity in all frames, it definitely implies the following properties:

i) any dispersion law describing particles generated by the field is Lorentz invariant, namely the corresponding hypersurface $\mathcal{M}$ is a sheet of hyperboloid with equation of the form $p_0 = \sqrt{\vec{p}^2 + m^2}$ (or the light cone ∂V^+ if $m = 0$).

ii) In all the sectors (or collision channels) of the space of states of the (interacting) field theory considered, the hypersurfaces which border the continuous part of the energy-momentum spectrum, including possible holes in the latter, are also Lorentz-invariant, namely sheets of hyperboloid of the form $p_0 = \sqrt{\vec{p}^2 + M_i^2}$ (or the light-cone).

It is the purpose of the present paper to give a hopefully elementary and self-contained proof of the latter facts, which have been established long ago in a general, although slightly different, framework by Borchers and Buchholz. As a matter of fact, the interest for the possible occurrence of Lorentz-symmetry breaking is not new and it has already been a subject of deep investigation in the framework of the basic principles of Quantum Field Theory (QFT): the latter two properties of Lorentz-invariance of the energy-momentum spectrum have indeed been proven in a paper by H.J. Borchers and D. Buchholz entitled “The Energy-Momentum Spectrum in Local Field Theories with Broken Lorentz-Symmetry” [2] completed by a paper by H.J. Borchers entitled “Locality and covariance of the spectrum” [3] in the general framework of Algebraic QFT (or “Local Quantum Physics”) [4]. In this deep analysis, generalizing similar results already obtained in [5] (see also [6] for a complete survey of the question), it was proven that the interplay of a weak form of microcausality, namely the commutativity of local observables attached to pairs of mutually space-like regions, together with energy-positivity in all Lorentz frames was sufficient to produce a Lorentz-invariant shape of the energy-momentum spectrum, even if the Lorentz-symmetry was broken in the considered physical representation of the field observables. In view of the always vivid interest of the community for the possible occurrence of some form of Lorentz-symmetry violation emerging from the spontaneous breaking at Planck scale of a “fundamental field or string theory” (see [1] and references therein), but also

of its apparent unawareness of the results of [2,3], we think it useful to give a revival to these results in a way which we hope to be accessible to the current field-theorist reader. In fact, we wish to give here a new presentation of these unexpected properties of geometrical nature in energy-momentum space in terms of propagators and Green's functions of microcausal Bose and Fermi-fields. We shall thus avoid using the less familiar formulation of Algebraic QFT, and will focus on the contrary on the phenomena of complex geometry which play a basic role in this matter and still hold in a context of indefinite-metric (see our concluding remarks).

Some preliminary comments on the exact content of properties i) and ii) (announced above) and on the relevant concepts and field-theoretical framework used in our approach are necessary.

As in [2,3], the proof of properties i) and ii) which we shall give is of a general nature, i.e. non-perturbative and in fact independent of any Lagrangian formulation of the field model. However, our method is close to the preoccupations of [1] by its use of such basic objects as the transforms of causal (retarded and advanced) propagators of the fields in energy-momentum space. In the approach of [1], the dispersion laws of particles are always associated with given quadratic parts of field Lagrangians incorporating explicit Lorentz-symmetry breaking coefficients of appropriate type. Such dispersion laws therefore correspond to particles which are “elementary” with respect to the field introduced in the Lagrangian, namely they appear as associated with poles of the propagator of this field in energy-momentum space. To be complete, another case of dispersion laws also deserves to be considered, namely those which correspond to “composite” particles of the field: the latter appear as associated with poles of the four-point (or higher n -point) functions of the field in energy-momentum space; for example, this is the case for the hadronic particles if the fundamental fields are those of the standard model.

Here we shall show in detail the previously announced geometrical properties of Lorentz invariance for the poles of propagators (corresponding to the case considered in [1]) and we shall also indicate the derivation of the corresponding equally valid results for the poles of four-point (or n -point) functions. We shall thus be concerned with stable (elementary or composite) particles, corresponding to discrete parts of the spectrum, not embedded in the continuum. The case of unstable particles corresponding to possible complex poles of the Green's functions in unphysical sheets is excluded from our study.

We wish to stress that the somewhat surprising phenomenon of geometrical Lorentz-invariance produced in the present problem has to do with peculiar properties of complex geometry in several complex variables. This phenomenon occurs as soon as the retarded and advanced propagators (or more general n -point functions) have Fourier-Laplace transforms, holomorphic in specific “tube domains” of complex energy-momentum space which correspond in a very general way to the causal support properties of these functions in spacetime, expressing the microcausality of the fields. ⁽²⁾ The traditional framework of QFT in which these *analyticity properties of Green's functions in complex energy-momentum space* have been derived (see [8,9,17] and references therein) is the Wightman-

⁽²⁾ Such properties, which are also closely related to the Jost-Lehmann-Dyson (JLD) formula for causal commutators[7], have been thoroughly exploited in [2,3] precisely in the spirit of [7].

LSZ [18,19] axiomatic framework in which the various n -point functions of the fields are supposed to be tempered distributions. As an application of the theory of Fourier-Laplace transforms of tempered distributions with causal supports, making use of the notion of “boundary values of analytic functions in the sense of distributions” (see e.g. [12]), the previously mentioned analyticity domains of the Green’s functions in momentum space have been obtained *together with polynomial increase properties of the functions in their respective domains*. From the viewpoint of the Lagrangian formalism of QFT, this framework is supposed to fit with *conventional (renormalizable) field models* whose Lagrangian only involves a finite number of derivatives of the fields. In the various examples of conventional type of [1] in which Lorentz- symmetry-breaking terms have been introduced so as to produce a non-Lorentz-invariant dispersion law (condition a)), it was found that the conditions b),c) listed at the beginning of this introduction could not be satisfied simultaneously: this fact is indeed consistent with our property i) since (as shown below) the violation of the group-velocity condition c) implies the violation of microcausality.

We now emphasize the existence of enlarged frameworks of QFT, making use of the general mathematical setting of hyperfunction theory (see e.g. [20] and references therein, the first generalization in this direction being the theory of Jaffe fields [21]). In the hyperfunction setting, the retarded and advanced Green’s functions in spacetime may have arbitrarily wild singularities; as we shall see below, this does not prevent their Fourier-Laplace transforms from being well-defined in the relevant complex domains of momentum space, although they no longer have a polynomial rate of increase at infinity in these domains. The analyticity properties of these transforms still express (at a high degree of generality) the microcausality property of such possible theories, which is now defined in terms of appropriately chosen classes of test-functions. It will then be clear from our geometrical proof of properties i) and ii) that this proof works for the class of *all holomorphic functions* in the domains considered, and therefore that the loss of polynomial increase properties in such general field theories does not invalidate the results. From the viewpoint of the Lagrangian formalism of QFT, this very general framework of QFT (implying no limitation on the strength of short-distance singularities) should cover (provided they exist!) all *nonconventional models* whose Lagrangian involves an infinite number of derivatives of the fields.

In our section 2, we shall first specify the basic analyticity properties of retarded and advanced two-point functions in tube domains, which express microcausality in complexified energy-momentum space; for completeness, we shall explain the result in elementary terms for the non-trivial case of QFT in the hyperfunction setting, in comparison with the standard result for Wightman fields in the setting of tempered distributions. This sketch of the general case is presented here (in Sec. 2.1) for convincing the reader that the validity of analyticity in complex momentum space is not submitted to restrictions due to lack of convergence of the Fourier integrals; however, this matter is outside our main purpose which is the derivation of properties i) and ii) and it may be skipped without inconvenience by the reader. Then we describe the procedure through which information on the energy-momentum spectrum is encoded in this general approach in complex momentum space. We then formulate three basic results of complex geometry, called Properties A, B and C, whose physical consequences in terms of *admissible dispersion laws* are derived

in a straightforward way: Property A explains why the velocity group condition c) of dispersion laws is implied by microcausality under a weak requirement of energy-positivity. Properties B and C provide a proof of the previous statements (i) and ii)) for the dispersion laws of elementary particles and for the thresholds (and possible holes) of the continuous spectrum, under the joint requirement of microcausality and energy-positivity in all frames. A complete proof of Properties A, B and C is given in this section. In section 3, it is shown that similar consequences of microcausality and (weak or strong) energy-positivity requirements can be formulated in terms of momentum-space analyticity properties of four-point (resp. more generally $2n$ -point) Green's functions established in [8,9] (resp. [17d),e])). The exact counterparts of Properties A,B,C, called respectively A',B',C', are then described and these phenomena of complex geometry are shown to imply the corresponding statements (i) and ii)) for the dispersion laws of composite particles and for the thresholds (and possible holes) of the continuous spectrum in the channels considered. Section 4 gives concluding remarks.

2 Shape of the energy-momentum spectral supports for the two-point functions

2.1 Expressing microcausality in complex momentum space

For simplicity, we first recall the *basic analyticity property in complex momentum space* by using conventional Fourier integrals (as if the fields were “good functions ” of x) before explaining why this property still holds with a high degree of generality in the rigorous settings of tempered distributions and hyperfunctions, corresponding respectively to conventional and nonconventional field theories.

Let $F^+(p)$ and $F^-(p)$ (with $p = (p_0, \vec{p})$) be respectively the Fourier transforms of the vacuum expectation values of the retarded and advanced (anti-)commutators of a general (fermion or boson) quantum field $\Phi(x)$:

$$F^+(p) = i \int e^{ip \cdot x} \theta(x_0) < [\Phi(\frac{x}{2}), \Phi(-\frac{x}{2})]_{\pm} > dx_0 d\vec{x}, \quad (1)$$

$$F^-(p) = -i \int e^{ip \cdot x} \theta(-x_0) < [\Phi(\frac{x}{2}), \Phi(-\frac{x}{2})]_{\pm} > dx_0 d\vec{x}. \quad (1')$$

For writing the latter, we have assumed as usual that the space of states in which the field is acting carries a representation of the group of spacetime translations and that the field is invariant under this representation; energy and momentum operators are the corresponding generators of this group. It is of current use to exploit the analyticity properties of $F^+(p)$ and $F^-(p)$ respectively in the upper and lower half-planes of the complexified energy variable p_0 . However, the postulate of microcausality for the field $\Phi(x)$ implies much more. In fact, it requires that the retarded and advanced propagators occurring under the integrals at the r.h.s. of Eqs (1) and (1') have respectively their supports contained in the closed forward and backward cones $\overline{V}^+ (x_0 \geq |\vec{x}|)$ and $\overline{V}^- (x_0 \leq -|\vec{x}|)$. It then follows that the integrals (1) and (1') remain convergent and define analytic functions of the complex energy-momentum vector $k = p + iq$, still denoted by $F^+(k)$ and $F^-(k)$,

in the respective domains T^+ (p arbitrary, $q \in V^+$) and T^- (p arbitrary, $q \in V^-$); $V^+ = -V^-$ is the open forward cone: $q_0 > |\vec{q}|$. T^+ and T^- are called the “forward and backward tubes”; they contain respectively the upper and lower half-planes in all their one-dimensional sections by (complexified) time-like straight lines, interpreted as energy variables in all possible Lorentz frames. $F^\pm(k)$ are the “Fourier-Laplace transforms” of the retarded and advanced propagators in complex energy-momentum space; their boundary values $F^\pm(p)$ on the reals from (respectively) $T^\pm$ are the Fourier transforms themselves of these propagators.

In view of the local singularities of the fields, the previous derivation of this basic analyticity property has to be corrected under two respects in realistic cases, namely i) splitting in a meaningful way the “(anti-)commutator function” $C(x) = \langle [\Phi(\frac{x}{2}), \Phi(-\frac{x}{2})]_\pm \rangle$ with support $\bar{V}^+ \cup \bar{V}^-$ into its retarded and advanced components $R(x)$ and $A(x)$ with respective supports $\bar{V}^+$ and $\bar{V}^-$, in such a way that $C(x) = -i(R(x) - A(x))$ (this was done formally in Eqs (1) and (1') by using the Heaviside multipliers $\theta(\pm x_0)$), and ii) giving a sense to the formal integrals of (1) and (1') when p is replaced by $k = p + iq$ respectively taken in T^+ and T^- .

In the framework of Wightman-LSZ fields where $C(x)$ is a tempered distribution, there exists a standard splitting procedure (see e.g. [9]) which defines $R(x)$ and $A(x)$ as tempered distributions with respective supports $\bar{V}^+$ and $\bar{V}^-$ (up to an ambiguity which is simply a linear combination of $\delta(x)$ with derivatives of the latter). Then in view of this support property the Fourier-Laplace exponential $e^{i(p+iq)\cdot x}$ is seen to be a good test-function, respectively for $R(x)$ provided q is taken in V^+ and for $A(x)$ provided q is taken in V^- , which justifies the corresponding analyticity property of $F^\pm(p+iq)$ in $T^\pm$. Moreover these functions are proved to have polynomial increase for $k = p+iq$ tending to infinity in $T^\pm$ and to be bounded by an inverse power of $|q|$ near the reals: for p real, $F^\pm(p)$ are then defined as distribution boundary values of $F^\pm(k)$ from the respective tube domains $T^\pm$. The Fourier transform $\tilde{C}(p)$ of $C(x)$ in the (L. Schwartz) sense of tempered distributions, which has to be a measure (for spectral reasons, see below), is then given by the following relation between distributions $\tilde{C}(p) = -i(F^+(p) - F^-(p))$.

In the enlarged framework of QFT where the definition of field operators is given in the sense of hyperfunction theory (see e.g. [20]), $C(x)$ is a hyperfunction on x -space, namely a functional on an appropriate space $\mathcal{A}$ of analytic functions $\varphi(x)$. However, if one needs this concept in order to include arbitrarily wild local singularities in x -space, one may keep as a basic requirement the fact that in momentum space, the “spectral function” $\tilde{C}(p) = \langle \tilde{\Phi}(p)\tilde{\Phi}(-p) \rangle - \langle \tilde{\Phi}(-p), \tilde{\Phi}(p) \rangle$ ⁽³⁾ remains a distribution and even a measure in p_0 , depending continuously on $\vec{p}$. This is still a consequence of the existence of a unitary representation of the spacetime translations acting on the field operators in the Hilbert space of states; more intuitively speaking, if one excludes phenomena of infrared divergences, the contributions $\delta(p_0 - \omega(\vec{p}))$ associated with dispersion laws of the theory are still considered as dominant singularities in momentum space. However, $\tilde{C}(p)$

⁽³⁾ Here the “bracket notation” in terms of operator products is used only for its suggestive content; no (infinite!) energy-momentum conservation δ -function is involved in it.

is no longer a *tempered* distribution (i.e. it may have nonpolynomial increase at infinity, in correspondence with the wild local singular behaviour of $C(x)$). Mathematically, this means that a reasonable space $\mathcal{A}$ of test-functions is the space of functions $\varphi(x)$ whose Fourier transforms $\tilde{\varphi}(p)$ belong to the Schwartz space $\mathcal{D}$ (functions infinitely differentiable with compact support): $\mathcal{A}$ is a space of entire functions $\varphi(x + iy)$ which decrease more rapidly than any inverse power of $|x|$ in the reals and are bounded by linear exponentials of y . Then the Fourier transformation relating C and $\tilde{C}$ is defined in the usual way by the formula $(C, \overline{\varphi}) = (\tilde{C}, \tilde{\overline{\varphi}})$. Typically, a hyperfunction $C(x)$ satisfying the previous requirements admits a “concrete representative” (defined up to an arbitrary entire function of x) which is a function $\hat{C}(z_0, \vec{x})$, holomorphic with respect to $z_0 = x_0 + iy_0$ in the cut-plane excluding the reals. $\hat{C}$ must be polynomially bounded at infinity (corresponding to the distribution character of $\tilde{C}(p)$), but its behaviour near the reals may be arbitrarily wild, corresponding to the idea that the hyperfunction $C(x)$ is (in a heuristic sense) the “jump of $\hat{C}$ across the reals”. What remains indeed concretely defined is the following integral (independent of η , $\eta > 0$, for all φ in $\mathcal{A}$):

$$(C, \varphi) = \int d\vec{x} \int_{-\infty}^{+\infty} dx_0 \left[\hat{C}(x_0 + i\eta, \vec{x}) \varphi(x_0 + i\eta, \vec{x}) - \hat{C}(x_0 - i\eta, \vec{x}) \varphi(x_0 - i\eta, \vec{x}) \right]. \quad (2)$$

Microcausality and splitting: the formulation of a support condition like microcausality cannot be done as in the case of distributions by using localized test-functions since $\mathcal{A}$ does not contain such functions. But making use of the representative $\hat{C}$ of C , one can formulate the microcausality condition by postulating that the jump of $\hat{C}$ across the reals vanishes in the complement of $\overline{V}^+ \cup \overline{V}^-$, or in other words that $\hat{C}(x_0, \vec{x})$ is holomorphic there (in x_0). Now it can be shown that a splitting of the form $\hat{C}(z_0, \vec{x}) = -i(\hat{R}(z_0, \vec{x}) - \hat{A}(z_0, \vec{x}))$ can always be done (with an ambiguity consisting in functions carrying a singularity only at $(x_0, \vec{x}) = 0$) in such a way that $\hat{R}$ and $\hat{A}$ be holomorphic in the cut z_0 -plane (for all $\vec{x}$), polynomially bounded at infinity and such that the jumps of $\hat{R}$ and $\hat{A}$ across the reals vanish respectively in the complements of $\overline{V}^+$ and $\overline{V}^-$. The feasibility of this splitting which is not surprising for $|\vec{x}| \neq 0$, (since the “retarded and advanced cuts” are separated in that case) requires more mathematical knowledge for $\vec{x} = 0$; such a decomposition belongs to a category called “Cousin’s problem” which is standard in complex analysis. Applying the procedure of formula (2) for defining the hyperfunctions $R(x)$ and $A(x)$ with respective representatives $\hat{R}$ and $\hat{A}$, one thus obtains for C a satisfactory splitting of the form $C(x) = -i(R(x) - A(x))$.

Analyticity in the tubes $T^\pm$: applying the definition (2) to R and A , and taking into account the location of the corresponding cuts for $\hat{R}$ and $\hat{A}$ in the corresponding formulae, one can perform a contour distortion argument which yields the following alternative forms:

$$(R, \varphi) = \int d\vec{x} \int_{\gamma^+(\vec{x})} dz_0 \hat{R}(z_0, \vec{x}) \varphi(z_0, \vec{x}), \quad (2')$$

$$(A, \varphi) = \int d\vec{x} \int_{\gamma^-(\vec{x})} dz_0 \hat{A}(z_0, \vec{x}) \varphi(z_0, \vec{x}), \quad (2'')$$

where $\gamma^+(\vec{x})$ and $\gamma^-(\vec{x})$ are folded contours which surround respectively the half-lines $x_0 \geq |\vec{x}|$ and $x_0 \leq -|\vec{x}|$ in the z_0 -plane. One now notices that the entire function $e^{i(p_0+iq_0)z_0-i(\vec{p}+i\vec{q})\cdot\vec{x}}$ is an acceptable test-function φ in (2') if $(q_0, \vec{q})$ is in V^+ , and in (2'') if $(q_0, \vec{q})$ is in V^- . One thus justifies again in this very general case the fact that $F^+(k) = (R, e^{ik\cdot x})$ and $F^-(k) = (A, e^{ik\cdot x})$ are respectively holomorphic in the tubes T^+ and T^- . Moreover, by taking into account the distance η of the contours $\gamma^\pm(\vec{x})$ to the corresponding cuts in the z_0 -plane, one would show that these Fourier-Laplace transforms of R and A now satisfy bounds of the following form in their respective tube domains:

$$|F^\pm(k)| \leq M_\eta e^{\eta(|p_0|+|q_0|)},$$

where the positive number η can be given an arbitrary value (the dependence on η of the corresponding constant M_η being unspecified).

2.2 A problem of analytic completion

So in the sector generated by “one-field vector-states” of the form $\int \varphi(x)\Phi(x)dx$, with φ varying in the relevant space of test-functions (corresponding to the field-theoretical framework which one wishes to consider), microcausality is fully expressed by the analyticity of the pair of functions (F^+, F^-) in the corresponding domains T^+, T^- . Now any usable information on the support of the energy-momentum spectrum of the theory in this sector amounts to specifying an open subset $\mathcal{R}$ of the (real) energy-momentum space in which the distributions $\langle \tilde{\Phi}(p)\tilde{\Phi}(-p) \rangle$ and $\langle \tilde{\Phi}(-p), \tilde{\Phi}(p) \rangle$ vanish together. In fact, such a support property implies the *coincidence relation* $F|_{\mathcal{R}}^+ = F|_{\mathcal{R}}^-$, since the expression

$$F^+(p) - F^-(p) = i\tilde{C}(p) = i \langle [\tilde{\Phi}(p), \tilde{\Phi}(-p)]_\pm \rangle \quad (3)$$

vanishes in $\mathcal{R}$. It then follows from a standard theorem of complex analysis called the “edge-of-the-wedge theorem” (still valid in the case of holomorphic functions with distribution-like boundary values, see [10] and references therein), that $F^+(k)$ and $F^-(k)$ then admit a *common analytic continuation* $F(k)$ which is analytic in the union of T^+, T^- and of a complex neighborhood of the real set $\mathcal{R}$; in other words, F^+ and F^- “communicate analytically” through $\mathcal{R}$, as functions of the set of complex variables $k = (k_0, \vec{k})$.

It is one of the basic phenomena of Analysis and Geometry in several complex variables that arbitrary (connected) subsets of complex space $\mathbf{C}^n$ are not in general “natural” for the class of holomorphic functions: this means that for such a general subset Σ , all the functions holomorphic in Σ admit an analytic continuation in a common larger domain $\hat{\Sigma}$, called the holomorphy envelope of Σ . This phenomenon, which does not exist in the single-variable case, involves exclusively geometrical properties of the set Σ and the extension from Σ to $\hat{\Sigma}$ can always be done in principle by an appropriate use of the Cauchy integral formula; this *analytic completion procedure* presents a strong analogy with the procedure of taking the convex hull $\check{S}$ of a subset S in the ordinary real space $\mathbf{R}^n$, the notion of a “natural holomorphy domain” in $\mathbf{C}^n$ being a certain generalisation of the notion of “convex domain” in $\mathbf{R}^n$ (see e.g. [11,12] and references therein). As a matter of fact, the most

standard and useful result in this connection is the so-called “tube theorem” (see e.g. [12]) which we shall apply below: *Any domain D in $\mathbf{C}^n$ which is “tube-shaped”, i.e. of the form $\mathbf{R}^n + iB$ admits a holomorphy envelope which is the tube $\hat{D} = \mathbf{R}^n + i\hat{B}$, where $\hat{B}$ is the convex hull of B in $\mathbf{R}^n$.*

It turns out that sets of the form $\Sigma_{\mathcal{R}} = T^+ \cup T^- \cup \mathcal{R}$ are not natural and that, for various choices of $\mathcal{R}$ of physical interest, the corresponding holomorphy envelope $\hat{\Sigma}$ or parts of it can be computed and unexpectedly strong results then follow. Cases when $\mathcal{R}$ itself can be extended to a larger real region $\hat{\mathcal{R}}$ (namely $\hat{\mathcal{R}} = \hat{\Sigma} \cap \mathbf{R}^n \supset \mathcal{R}$) are specially interesting, since they correspond to enlarging the region on which the “spectral function” $\langle [\tilde{\Phi}(p), \Phi(-p)]_{\pm} \rangle$ is proven to vanish, and therefore to refining our information on the support of the distribution $\langle \tilde{\Phi}(p)\tilde{\Phi}(-p) \rangle$, called “spectral support”. Properties A and C given below are precisely of this type. Property B is a basic example of a holomorphy envelope for a domain $\Sigma_{\mathcal{R}}$ which exactly corresponds to the case when energy-positivity is satisfied in all frames.

2.3 Microcausality implies dispersion laws with subluminal velocities

If energy-positivity is required to hold *only in privileged frames, such as the laboratory frame and a set of frames which have small velocities with respect to the latter* ⁽⁴⁾, there exists a maximal region $\hat{\mathcal{R}}$ of the form $-\omega(\vec{p}) < p_0 < \omega(\vec{p})$ (with $\omega(\vec{p}) \geq \gamma|\vec{p}|$ for some positive constant γ) in which the (anti-)commutator function $\langle [\tilde{\Phi}(p), \tilde{\Phi}(-p)]_{\pm} \rangle$ vanishes. We claim that, *due to microcausality, the hypersurface with equation $p_0 = \omega(\vec{p})$ is not arbitrary: it has to be a space-like hypersurface.* In fact, the geometry of the relativistic light-cone is deeply involved in the implications of microcausality, as it results from the following

Property A (“Double-cone theorem”):

Let $\mathcal{R}_{a,b}$ be a neighborhood (in real p -space) of a given time-like segment $]a, b[$ with end-points a and b (b in the future of a). Then any function $F(k)$ holomorphic in $\Sigma_{\mathcal{R}_{a,b}}$ admits an analytic continuation in a (complex) domain which contains the real region $\diamond_a^b$, where $\diamond_a^b$ is the “double-cone” defined as the set of all points p such that p is in the future of a and in the past of b .

Interpretation of Property A:

Let $\mathcal{M}$ ($p_0 = \omega(\vec{p})$) be the hypersurface bordering the vanishing region $\hat{\mathcal{R}}$ of the (anti-)commutator function of a certain field theory satisfying microcausality and energy-positivity in privileged frames. Then for each point $b = (\omega(\vec{p}), \vec{p})$ in $\mathcal{M}$, there exists some interval of the form $\omega(\vec{p}) - \epsilon < p_0 < \omega(\vec{p})$ and some open neighborhood $\mathcal{R}_{a,b}$ of the time-like segment $]a, b[$ defined by this interval (i.e. $a \equiv (\omega(\vec{p}) - \epsilon, \vec{p})$) which lies in $\hat{\mathcal{R}}$. It then follows from Property A that the propagator $F(k)$ of this theory has to be analytic in the full double-cone $\diamond_a^b$, and therefore that the corresponding (anti-)commutator function must vanish in this double-cone: therefore, $\diamond_a^b$ belongs to $\hat{\mathcal{R}}$, and this argument holds for every point b of $\mathcal{M}$, which shows that $\mathcal{M}$ has to be a spacelike hypersurface.

Similarly, assume that the vanishing region $\hat{\mathcal{R}}$ is accompanied by another pair of maximal vanishing regions $\hat{\mathcal{R}}_1^{\pm}$ of the form $\omega(\vec{p}) < |p_0| < \omega_1(\vec{p})$ of the (anti-)commutator

⁽⁴⁾ This refers to the notion of “concordant frames” introduced in [1]

function. Then $p_0 = \omega(\vec{p})$ appears as the dispersion law of a particle corresponding to a pole $\frac{Z(\vec{p})}{k_0 - \omega(\vec{p})}$ of the propagator $F(k)$. So the previous argument shows in this case that both the hypersurface $\mathcal{M}$ describing the dispersion law of the particle and the hypersurface $\mathcal{M}_1$ ($p_0 = \omega_1(\vec{p})$) bordering the region $\hat{\mathcal{R}}_1^+$ have to be spacelike. The argument extends of course to the case of any (ordered) set of dispersion laws corresponding to several particles. *Therefore, for every particle appearing with an energy gap in the propagator of the field considered, microcausality alone implies that condition c) (subluminal or luminal velocities) is satisfied by such a particle.*




Fig. 1. The “double-cone” $\diamond$ and the curves $\check{h}_\lambda$

Proof of Property A:

This theorem, which can be seen as a generalisation of a similar property (corollary of the “mean value Asgeirsson theorem”) for the solutions of the wave-equation [13], has been proved by Vladimirov [14] and by Borchers [15]. The main geometrical idea is displayed by treating a typical case in two-dimensional energy-momentum space with coordinates (p_0, p_1) . We take for $]a, b[$ the segment $\delta =]-1, +1[$ of the time axis and for $\mathcal{R}_{a,b}$ a thin rectangle δ_ϵ of the form: $|p_0| < 1$, $|p_1| < \epsilon$. The tubes T^+, T^- in the complexified space with coordinates $(k_0 = p_0 + iq_0, k_1 = p_1 + iq_1)$ are defined respectively by the conditions $q_0 + q_1 > 0$, $q_0 - q_1 > 0$ and $q_0 + q_1 < 0$, $q_0 - q_1 < 0$, and we shall show that the real region obtained by analytic completion of $T^+ \cup T^- \cup \delta_\epsilon$ contains the “double-cone” $\diamond$ (a square in this case!) defined by the inequalities: $|p_0 - p_1| < 1$, $|p_0 + p_1| < 1$. One introduces the

family of complex curves h_λ with equation $[k_0^2 - (k_1 - 1)^2] = \lambda[k_0^2 - (k_1 + 1)^2]$, where the parameter λ varies in a complex neighborhood $\mathcal{V}$ of the real interval $]0, +\infty[$. Except for h_1 which is the (complexified) p_0 -axis, all these curves are hyperbolae, and $\diamond$ is generated by the (real) arcs $\check{h}_\lambda$ of h_λ parametrized by $-1 < p_0 < 1$ (with $|p_1| < 1$) when λ varies from 0 to $+\infty$; in a subinterval of the form $|\lambda - 1| < \eta$ (for some η determined by ϵ), $\check{h}_\lambda$ remains inside the rectangle δ_ϵ (see fig 1).

One then checks that for any function $F(k_0, k_1)$ holomorphic in $T^+ \cup T^- \cup \delta_\epsilon$ the change of complex variables $(k_0, k_1) \rightarrow (k_0, \lambda)$ is admissible. It allows one to define $\underline{F}(k_0, \lambda) = F(k_0, k_1(k_0, \lambda))$ as an analytic function for λ varying in $\mathcal{V}$ and k_0 varying in a ring-shaped domain D_λ , which surrounds and *excludes a neighborhood of a real interval of the form* $-1 + \alpha(\lambda) \leq p_0 \leq 1 - \alpha(\lambda)$ (fig 2a). This comes from the fact that for $0 < \lambda < +\infty$, the full upper (resp. lower) half-plane in the variable k_0 represents a set of points (k_0, k_1) of h_λ in T^+ (resp. T^-) ⁽⁵⁾ and that these two half-planes are connected by small real intervals $] -1, -1 + \alpha[$, $]1 - \alpha, 1[$ which represent points in δ_ϵ . Moreover, for $1 - \eta < \lambda < 1 + \eta$ *the full unit disk* $|k_0| < 1$ is in the analyticity domain of $\underline{F}$ (fig 2b), the corresponding arcs $\check{h}_\lambda$ being all contained in δ_ϵ .




Fig. 2. Initial analyticity domains of $\underline{F}(k_0, \lambda)$ in the k_0 -plane

⁽⁵⁾ To see this, one can e.g. rewrite the equation of h_λ as follows: $\frac{U-1}{U+1} = \lambda \frac{V-1}{V+1}$ with $U = k_0 + k_1$, $V = k_0 - k_1$, which entails (for $\lambda > 0$) the condition $\Im m U \times \Im m V > 0$, and therefore the fact that all complex points $(k_0, k_1) \equiv (U, V)$ in h_λ belong either to T^+ or to T^- according to whether $\Im m k_0 \equiv \frac{1}{2}(\Im m U + \Im m V)$ is positive or negative.

Now consider the Cauchy integral

$$I(k_0, \lambda) = \frac{1}{2i\pi} \oint_{\gamma_\lambda} \frac{\underline{F}(k'_0, \lambda)}{k'_0 - k_0} dk'_0,$$

where γ_λ has its support in D_λ , is homotopic to the (anticlockwise) unit circle and equal to it for $\lambda = 1$. In view of the latter analyticity property of $\underline{F}$, one has $I(k_0, \lambda) = \underline{F}(k_0, \lambda)$ for $1 - \eta < \lambda < 1 + \eta$ and therefore $I(k_0, \lambda)$ provides an analytic continuation of $\underline{F}(k_0, \lambda)$ itself inside the full disk bordered by γ_λ and therefore on the real interval $] -1, +1[$ of the variable k_0 , for all λ in the interval $]0, +\infty[$. By coming back to the original variables (k_0, k_1) , this shows that F admits an analytic continuation in the full region $\diamond$. In the most general version of the theorem in two dimensions, the neighborhood $\mathcal{R}_{a,b}$ of the given time-like segment $]a, b[$ is considered as a union of rectangles of the previous δ_ϵ -type, whose thickness ϵ tends to zero while they tend to $]a, b[$: the double-cone (or square) $\diamond_{a|d=2}^b$ is then clearly obtained as the union of the corresponding squares $\diamond$ obtained in the previous procedure of analytic completion. Finally the proof of the theorem in the d -dimensional case is obtained by applying the two-dimensional result in all the planar sections passing by a and b , since i) the two-dimensional sections of the tubes $T^\pm$ are the corresponding tubes of the (complexified) planar sections, and ii) $\diamond_a^b$ is generated by the union of all double-cones of the previous type $\diamond_{a|d=2}^b$ in these planar sections.

2.4 Microcausality and energy-positivity in all frames imply Lorentz invariant spectral supports

A basic implication of microcausality together with energy-positivity in all frames is the fact that propagators $F(k)$ of the underlying fields have to be *holomorphic in a domain which is invariant under all complex Lorentz transformations*, even if these propagators are not Lorentz invariant functions due to the fact that the Lorentz symmetry is broken in the representation of the fields under consideration. The key property which is at the origin of this peculiarity is the following

Property B (“Källen-Lehmann domain”):

Let $\mathcal{R} = \mathcal{R}_0$ be the set of all space-like energy-momentum vectors $p = (p_0, \vec{p}) : |p_0| < |\vec{p}|$. Then any function $F(k)$ holomorphic in $\Sigma_{\mathcal{R}_0} = T^+ \cup T^- \cup \mathcal{R}_0$ admits an analytic continuation in the domain $\hat{\Sigma}_{\mathcal{R}_0}$ which is the set of all complex vectors $k = (k_0, \vec{k})$ such that $k^2 \equiv k_0^2 - \vec{k}^2$ is different from any positive number and from zero.

Interpretation of Property B:

Energy-positivity in all Lorentz frames implies that the distribution $\langle \tilde{\Phi}(p) \tilde{\Phi}(-p) \rangle$ vanishes in the complement of $\bar{V}^+$ and therefore, in view of (3), that the coincidence relation $F_{|\mathcal{R}_0}^+ = F_{|\mathcal{R}_0}^-$ is satisfied. Property B then implies the analyticity of the propagator $F(k)$ in the full “cut-domain” $\hat{\Sigma}_{\mathcal{R}_0}$. Our denomination of “Källen-Lehmann domain” for the latter is motivated by the fact that in the usual case when Lorentz invariance (or covariance) of the field is postulated, the analyticity domain $\hat{\Sigma}_{\mathcal{R}_0}$ is directly obtained as a byproduct of the Källen-Lehmann integral representation of the propagator

$$F(k) \equiv \underline{F}(k^2) = \int_0^\infty \frac{\rho(\sigma)}{\sigma - k^2} d\sigma,$$

since the image of $\hat{\Sigma}_{\mathcal{R}_0}$ in the variable k^2 is the usual cut-plane domain $\mathbf{C} \setminus \mathbf{R}^+$. Here, however, this Lorentz-invariant domain (considered in the full complex k -space) is obtained *without any assumption of Lorentz covariance and of boundedness of the functions*, but purely on the basis of microcausality and energy-positivity.

Moreover, one will show that any further information on the spectral support which is superimposed to the conditions of Property B implies the Lorentz-invariant shape of all the components of the spectral support together with the invariance under complex Lorentz transformations of the corresponding analyticity domain of the propagator. This is the purpose of the following property, whose statement in the present form is valid for any spacetime dimension $d \geq 3$; we postpone to the proof the corresponding statement for the two-dimensional case, which requires a little more care in view of the decomposition of the light-cone into two straight-lines (the so-called “left and right-movers”).

Property C (Lorentz-invariance of the borders of the spectral supports); case $d \geq 3$: If $\mathcal{R}$ is any real open set, not necessarily connected, containing $\mathcal{R}_0$ then every function $F(k)$ holomorphic in $\Sigma_{\mathcal{R}} = T^+ \cup T^- \cup \mathcal{R}$ admits an analytic continuation in the (Lorentz-invariant) set $\hat{\mathcal{R}}$ of all real vectors p whose Minkowskian norm p^2 has a value already taken at some vector in $\mathcal{R}$. Moreover the domain $\hat{\Sigma}_{\mathcal{R}}$ in which every such function $F(k)$ can be analytically continued is the set of vectors k such that k^2 takes all possible complex values and all real values taken by p^2 when p varies in $\mathcal{R}$.

Interpretation of Property C:

It is easy to see that Property C (in its first part) implies that if microcausality and energy-positivity are satisfied, then the most general type of set $\hat{\mathcal{R}}$ where the (anti-)commutator function (3) has to vanish is a set composed of one distinguished region $\mathcal{R}_{M_0}$ of the form $-\infty < p^2 < M_0^2$, with $M_0 \geq 0$ and of zero, one or several disjoint Lorentz-invariant regions of the form $M_i'^2 < p^2 < M_i^2$, where $M_1' \geq M_0$ and $M_{i+1}' \geq M_i$, $i = 1, \dots, l-1$, $M_l \leq \infty$. This implies in turn that the support of $\langle \tilde{\Phi}(p)\tilde{\Phi}(-p) \rangle$ is exactly the union of all the “thick (or thin) hyperbolic shells” defined by $M_i^2 \leq p^2 \leq M_{i+1}'^2$, $p_0 \geq 0$, ($i = 0, 1, \dots, l-1$), $p^2 \geq M_l^2$ (and of the origin if $\langle \tilde{\Phi}(p) \rangle \neq 0$). The equality case $M_i = M_{i+1}'$ corresponds to some “thin shell” $p^2 = M_i^2$. This thin shell situation occurs precisely when the distribution $\langle \tilde{\Phi}(p)\tilde{\Phi}(-p) \rangle$ describes a particle with dispersion law $p_0 = \sqrt{p^2 + M_i^2}$. No possibility is left for a Lorentz-symmetry breaking dispersion law. (Note that in this argument, the positivity of the Hilbert-space norm, implying the fact that the previous distribution is a positive measure factoring out a $\delta(p^2 - M_i^2)$, has not been used).

The proofs of Properties B and C given below are based on purely geometrical arguments. Both of them rely on a standard analytic completion procedure of geometrical type, namely the “tube theorem” (stated at the beginning of this section); apart from the recourse to this piece of knowledge in complex geometry, these proofs are completely self-contained. The analytic completion procedure is actually at work in the two-dimensional case, which we treat at first, while the general d -dimensional case will be reducible to the latter.

For the two-dimensional case, Property C must be properly restated as follows:

Property C (Lorentz-invariance of the borders of the spectral supports); case $d = 2$:

If $\mathcal{R}$ is any real open set, not necessarily connected, containing $\mathcal{R}_0$ then every function $F(k)$ holomorphic in $\Sigma_{\mathcal{R}} = T^+ \cup T^- \cup \mathcal{R}$ admits an analytic continuation in the (Lorentz-invariant) set $\hat{\Sigma}_{\mathcal{R}}$ obtained by adding to $\hat{\Sigma}_{\mathcal{R}_0}$ the set of all (real or complex) vectors k obtained by the action of real or complex Lorentz transformations on all vectors in $\mathcal{R}$.

We note that in the d -dimensional case, the latter version of Property C is equivalent to the former. In fact, the set of all vectors k obtained from a given vector $p = \underline{p} \neq 0$ in $\mathcal{R}$ by real or complex Lorentz transformations is the full complexified hyperboloid $k^2 = \underline{p}^2$ if $\underline{p}^2 \neq 0$ or the full complexified light cone $k^2 = 0$ if $\underline{p}^2 = 0$. However in the two-dimensional case, the latter statement differs from the former if $\mathcal{R}$ contains vectors $\underline{p}$ such that $\underline{p}^2 = 0$. In that case, the set of vectors k obtained from such a vector $\underline{p}$ by the action of real or complex Lorentz transformations is *not the full light cone* but only the complexified line of left or right-movers which the given vector $\underline{p}$ itself belongs to. In other words, one of these two lines may very well be a singular set of the propagator, and therefore contribute to the spectral support, although the other line doesn't; in such a case the parity symmetry of the spectral support is then broken but its Lorentz invariance is still preserved.

Proof of Properties B and C in the two-dimensional case:

We here consider the case when $k = (k_0, k_1)$ varies in $\mathbf{C}^2$, corresponding to two-dimensional field-theory. In the complex variables ($U = k_0 + k_1$, $V = k_0 - k_1$), the domains $T^{\pm}$ are described as $T^+ : \Im m U > 0, \Im m V > 0$, $T^- : \Im m U < 0, \Im m V < 0$, and $\mathcal{R}_0$ is the real set: $p^2 = UV < 0$. Let us then pass to the logarithmic variables $u = \log U$, $v = \log V$ and use the fact that any function $F(k) \equiv F(U, V) = F(e^u, e^v) \equiv f(u, v)$ is holomorphic and 2π -periodic with respect to the variables u and v in the image of $T^+ \cup T^- \cup \mathcal{R}_0$ in the space of these variables. One easily sees that the domain T^+ is one-to-one mapped (periodically) onto each one of the following (tube-shaped) domains $\Theta_l^+ = \mathbf{R}^2 + iB_l^+$ (l integer) where B_l^+ is the square $0 < \Im m u - 2l\pi < \pi$, $0 < \Im m v + 2l\pi < \pi$ and similarly for T^- onto each one of the domains $\Theta_l^- = \mathbf{R}^2 + iB_l^-$ (l integer) where B_l^- is the square $-\pi < \Im m u - 2l\pi < 0$, $\pi < \Im m v + 2l\pi < 2\pi$. As seen on fig 3, the set of all squares B_l^+ and B_l^- form a connected set *if one adds to them the common boundary vertices* represented by all the points $b_{\frac{l}{2}} = (\Im m u = l\pi, \Im m v = (-l+1)\pi)$, with l integer. But as one easily checks, the sets $\theta_{\frac{l}{2}} = \mathbf{R}^2 + ib_{\frac{l}{2}}$ belong precisely to the (periodic) image of the set $\mathcal{R}_0$ ($UV = e^{u+v} < 0$; e^u, e^v real). The function $f(u, v)$ is therefore holomorphic in the union of all the tube-shaped sets Θ_l^+ , Θ_l^- and $\theta_{\frac{l}{2}}$ and even (in view of the invariance of this edge-of-the-wedge configuration by all real translations in $\mathbf{R}^2$) in a *connected open tube* $\Theta = \mathbf{R}^2 + iB$ such that B is the union of all sets B_l^+ , B_l^- together with *open neighborhoods* of all the points $b_{\frac{l}{2}}$. Then in view of the tube theorem, $f(u, v)$ admits a (2π -periodic) analytic continuation in the tube $\hat{\Theta} = \mathbf{R}^2 + i\check{B}$, where $\check{B}$, namely the convex hull of B , is (as shown by fig 3) the domain $\check{B} : 0 < \Im m u + \Im m v < 2\pi$. $F(k)$ therefore admits an analytic continuation in the inverse image of the tube $\hat{\Theta}$ in the original variables, which is the set of all $k \equiv (U, V)$ such that $0 < \arg U + \arg V \equiv \arg k^2 < 2\pi$, namely the domain $\hat{\Sigma}_{\mathcal{R}_0}$ described in Property B.




Fig. 3. The set B (dark gray)
and its convex hull $\tilde{B}$ (light gray)

The domain $\hat{\Sigma}_{\mathcal{R}_0}$ can also be seen as the union of all complex hyperbolae h_ζ in $\mathbf{C}^2$ with equation $k^2 = UV = \zeta$ such that ζ belongs to the cut-plane $\mathbf{C} \setminus [0, \infty[$. Let us now assume that in addition to $\mathcal{R}_0$, the set $\mathcal{R}$ contains a given point $\underline{p} = (\underline{U}, \underline{V})$ with $\underline{p}^2 = \underline{\zeta} \geq 0$. To be specific, consider the case when one has: $\underline{U} > 0$ and $\underline{V} \geq 0$ and put $\underline{U} = e^{\underline{t}} > 0, \underline{V} = \zeta e^{-\underline{t}} \geq 0$, with $\underline{t}$ real; the remaining cases would be treated similarly by i) exchanging the roles of U and V and ii) changing (U, V) into $(-U, -V)$ in the following. We now use the fact that any function $F(k) \equiv F(U, V)$ analytic in $\Sigma_{\mathcal{R}} = T^+ \cup T^- \cup \mathcal{R}$ is analytic in a complex neighborhood of $\underline{p}$ and therefore in particular in a set of the form $\mathcal{N}(\underline{p}) = \{k = (U, V); U = e^t, V = \zeta e^{-t}; (\zeta, t) \in S_1\}$, where $S_1 = \{(\zeta, t); \underline{\zeta} - \epsilon < \zeta < \underline{\zeta} + \epsilon, |t - \underline{t}| < \rho\}$. It also follows from Property B that the image G of such a function $F(U, V)$ in the space of complex variables (ζ, t) , namely $G(\zeta, t) \equiv F(e^t, \zeta e^{-t})$, is analytic in the set $S_2 = \{(\zeta, t); |\zeta - \underline{\zeta}| < \epsilon, \Im \zeta \neq 0; t \in \mathbf{C}\}$ (with periodicity with respect to the translations $t \rightarrow t + 2il\pi$). Putting these two facts together, namely the analyticity of $G(\zeta, t)$ in the union of the sets S_1 and S_2 , and making

the new change of variables

$$\alpha = \log \frac{\zeta - \underline{\zeta} + \epsilon}{\underline{\zeta} + \epsilon - \zeta}, \quad \beta = i \log(t - \underline{t}),$$

one checks that the function $g(\alpha, \beta) \equiv G(\underline{\zeta} + \epsilon \frac{e^\alpha - 1}{e^\alpha + 1}, \underline{t} + e^{-i\beta})$ is holomorphic in the following tube-shaped domain $\mathcal{T} = \mathbf{R}^2 + i\mathcal{B}$, where $\mathcal{B}$ is the union of the (disconnected) open set $\{(\Im \alpha, \Im \beta); 0 < |\Im \alpha| < \frac{\pi}{2}; \Im \beta \text{ arbitrary}\}$ with the “connection interval” $\{(\Im \alpha, \Im \beta); \Im \alpha = 0; \Im \beta < \log \rho\}$ (see fig 4). Now since the convex hull of $\mathcal{B}$ is obviously the domain $\check{\mathcal{B}} = \{(\Im \alpha, \Im \beta); \frac{-\pi}{2} < \Im \alpha < \frac{\pi}{2}; \Im \beta \text{ arbitrary}\}$, the tube theorem implies that $g(\alpha, \beta)$ admits an analytic continuation in $\mathbf{R}^2 + i\check{\mathcal{B}}$, and therefore that $G(\zeta, t)$ admits an analytic continuation in the set $\{(\zeta, t); |\zeta - \underline{\zeta}| < \epsilon, t \in \mathbf{C}\}$ (with periodicity with respect to the translations $t \rightarrow t + 2il\pi$). Coming back to $F(U, V)$, this shows that F admits an analytic continuation in a set which is the union of all complex curves parametrized by $U = e^t, V = \zeta e^{-t}; t \in \mathbf{C}$, for ζ varying in the disk $|\zeta - \underline{\zeta}| < \epsilon$. These curves are complex hyperbolae except for the one corresponding to the value $\zeta = 0$, which is the straight-line $V = 0$, namely the (complexified) “right-mover” component of the light-cone. All these curves can be seen as generated by the action of all real or complex Lorentz transformations (parametrized by t) on the set $\mathcal{N}(\underline{p})$ and Property C is therefore established for the two-dimensional case.




Fig. 4. The set $\mathcal{B}$ (gray)

As a by-product of the latter, we stress the following result which is used below:

Property B with masses:

Let $\mathcal{R} = \mathcal{R}_\mu$ be the set of all real energy-momentum vectors p such that $p^2 < \mu^2$. Then any function $F(k)$ holomorphic in $\Sigma_{\mathcal{R}_\mu} = T^+ \cup T^- \cup \mathcal{R}_\mu$ admits an analytic continuation in the domain $\hat{\Sigma}_{\mathcal{R}_\mu}$ which is the set of all complex vectors k such that k^2 belongs to the cut-plane $\mathbf{C} \setminus [\mu^2, +\infty[$.

Remark It is sufficient that $\mathcal{R}$ is known to contain (neighborhoods of) one point $\underline{p}$ on the line $V = 0$ and one point $\underline{p}'$ on the line $U = 0$ (besides $\mathcal{R}_0$) in order to obtain an analyticity domain $\hat{\Sigma}_{\mathcal{R}}$ of the previous type $\hat{\Sigma}_{\mathcal{R}_\mu}$: in fact, Property C implies that both complex lines $U = 0$ and $V = 0$ are contained in the domain, except maybe for the point $U = V = 0$ which is not obtained by the previous analytic completion procedure. However, this point must also belong to the domain since an analytic function of two complex variables *cannot be singular at an isolated point* surrounded by its domain of analyticity (see e.g. [11]): it admits an analytic continuation at this isolated point defined by an appropriate Cauchy integral.

Proof of Properties B and C in the d -dimensional case:

The general case when $k = (k_0, \vec{k})$ varies in $\mathbf{C}^d$ (e.g. $d = 4$ for field theory in the physical Minkowskian space) will be treated by appropriately using the previous two-dimensional results in sections of $\mathbf{C}^d$ by (complexified) planes containing a time-direction.

Let $\underline{k} = \underline{p} + i\underline{q}$ be any vector in $\mathbf{C}^d$ such that $\underline{k}^2 \in \mathbf{C} \setminus [0, +\infty[$. In the affine Minkowskian space $\mathbf{R}^d$ consider the point P such that $[OP] \equiv \underline{p}$ and the time-like plane Π passing by P and generated by $\underline{q}$ and the unit vector e_0 of the time-axis (or choose one of these planes and call it Π in the degenerate case when $\underline{q}$ is along e_0 or is the null vector). There is a unique decomposition $\underline{p} = \underline{p}' + p_\perp$ such that $\underline{p}'$ is parallel to Π and $p_\perp$ is orthogonal to Π and therefore spacelike, if not the null vector: $p_\perp^2 = -\rho^2 \leq 0$. Introducing the complexified space $\Pi^{(c)}$ of Π and the two-dimensional vector variable $k' = p' + iq$ such that every point $k = p + iq$ in $\Pi^{(c)}$ can be uniquely written as $k = k' + p_\perp$ with k' orthogonal to $p_\perp$, one has: $k'^2 = k^2 + \rho^2$. In $\Pi^{(c)}$ the section of the domain $\Sigma_{\mathcal{R}_0} = T^+ \cup T^- \cup \mathcal{R}_0$ is represented in the vector-variable k' as the union of the two-dimensional tubes T'^+ and T'^- defined by $\Im m k'^2 > 0$ and respectively $\Im m k'_0 > 0$, $\Im m k'_0 < 0$, and of the real region defined by $p'^2 = p^2 - p_\perp^2 = p^2 + \rho^2 < \rho^2$. Therefore since the given vector $\underline{k} = \underline{p}' + p_\perp + i\underline{q} \equiv \underline{k}' + p_\perp$ is such that $\underline{k}'^2 = \underline{k}^2 + \rho^2 \in \mathbf{C} \setminus [\rho^2, +\infty[$, it follows from the two-dimensional *Property B with masses*, applied in k' -space to the restriction $F'(k') = F|_{\Pi^{(c)}}(k)$ of any function $F(k)$ analytic in $\Sigma_{\mathcal{R}_0}$, that F' admits an analytic continuation at $\underline{k}'$ and therefore that F itself can be analytically continued at the given vector $\underline{k}$. This shows that Property B holds in the d -dimensional case.

Proof of Property C: let us assume that in addition to $\mathcal{R}_0$, the set $\mathcal{R}$ contains a given vector $\underline{p} = [OP]$ with $\underline{p}^2 \geq 0$. Considering at first the case $\underline{p}^2 > 0$, we know that the two-sheeted hyperboloid $H(P)$ with equation $p^2 = \underline{p}^2$ can be seen as the union of all the hyperbolae $h_\alpha(P)$ passing by P which are the sections of $H(P)$ by all the two-dimensional planes Π_α containing the parallel to the time axis passing by P . In the complexified space of each (Minkowskian-type) plane Π_α , the domain $\Sigma_{\mathcal{R}}$ admits a restriction represented

by a two-dimensional domain of the form $\Sigma_{\mathcal{R}_\alpha}$, where $\mathcal{R}_\alpha$ contains P in addition to a region of the form $p_\alpha^2 < \rho_\alpha^2$, corresponding to the intersection of $\mathcal{R}_0$ by Π_α . Therefore, in view of Property C for the two-dimensional case the whole hyperbola $h_\alpha(P)$ (and even its complexified) belongs to the holomorphy envelope $\hat{\Sigma}_{\mathcal{R}_\alpha}$ of $\Sigma_{\mathcal{R}_\alpha}$. Since this is true for all hyperbolae $h_\alpha(P)$, the full hyperboloid $H(P)$ itself belongs to the holomorphy envelope $\hat{\Sigma}_{\mathcal{R}}$ of $\Sigma_{\mathcal{R}}$. In the case $\underline{p}^2 = 0$ (with $P \neq 0$), $H(P)$ is the light-cone and the previous argument of analytic completion in the union of all hyperbolic sections by the planes Π_α yields the whole light-cone *deprived from* the “light-ray” distinct from $[OP]$ and contained in the (unique) plane Π_{α_0} passing by the origin. However, this exceptional light-ray can be recovered by replacing P by a neighbouring point P' also such that $[OP']^2 = 0$: this is always possible since $\mathcal{R}$ is an open set. (We also note that for the same reason one thus obtains in that case an *open set* $\hat{\mathcal{R}}$ of the form $p^2 < \epsilon^2$, the isolated point $p = 0$ being also obtained according to the remark given at the end of the two-dimensional case). We have thus established the first part of Property C, namely the analytic completion at all *real* vectors p whose corresponding value of p^2 is taken by some vector $\underline{p} = [OP]$ in $\mathcal{R}$.

In order to establish the second part, we can now assume that $\mathcal{R}$ is the union of $\mathcal{R}_0$ together with a set of hypersurfaces H_μ of the form $p^2 = \mu^2$, with $\mu \geq 0$; then there remains to prove that all the points of the corresponding *complex* hypersurfaces $H_\mu^{(c)}$ can be reached by the previous analytic completion procedure. Here again, one can proceed as in the proof of Property B, namely taking any given vector $\underline{k} = \underline{p} + i\underline{q}$ in $H_\mu^{(c)}$, one considers the complex two-dimensional configuration in the corresponding plane $\Pi^{(c)}$ (specified above in the proof of Property B). Now the section of H_μ by the plane Π is a hyperbola contained in the region $\mathcal{R}_\Pi$ of the corresponding section, so that as a result of Property C in the two-dimensional case, the holomorphy envelope contains all the points of the corresponding *complex* hyperbola, which includes by construction the given point $\underline{k}$. For the case $\mu = 0$, the same method still works, including the treatment of the vectors $\underline{k} = \underline{p} + i\underline{q}$ such that $p^2 = q^2 = 0$, which belong to complexified light-rays: the latter are again obtained by the two-dimensional version of Property C in the special case of the right and left movers (no complex light-ray can be excluded since each light-ray has all its real points in the analyticity domain). This ends the proof of Property C in the general case.

3 Shape of the energy-momentum spectral supports for the N -point functions

We shall now show that the previous study can be repeated *for the sector generated by “two-field vector-states” of the form $\int \varphi(x, x') \Phi(x) \Phi(x') > dx dx'$* . It is in fact possible to perform a similar treatment in complex momentum space, in which propagators of the fields are now replaced by four-point functions of the latter: the corresponding results on the form of dispersion laws will then apply to composite particles appearing as “two-field bound-states”. Subsequently, we shall indicate the existence of a similar treatment for the sectors of “ n -field vector states” in terms of $2n$ -point Green’s functions with applications to dispersion laws of composite particles appearing as “ n -field bound states”, with $n \geq 3$. The validity of such a general study relies in an essential way on the general formalism of the analytic Green’s functions of interacting fields in complex momentum space [17].

The basic fact is that there exists an analog of formula (3) for the four-point function,

which can be written as follows (see again ⁽³⁾ for our use of the bracket notation):

$$F^+(p; p_1, p_2) - F^-(p; p_1, p_2) = < [\tilde{R}(p_1, p - p_1), \tilde{R}(p_2, -p - p_2)]_{\pm} > \quad (4)$$

where $\tilde{R}$ denotes the Fourier transform of a retarded two-point field operator carrying the total energy-momentum p :

$$\tilde{R}(p', p - p') = \int e^{ip \cdot x} e^{ip' \cdot (x' - x)} \theta(x'_0 - x_0) [\Phi(x'), \Phi(x)]_{\pm} dx dx' \quad (5)$$

and where $F^+(p; p_1, p_2)$ and $F^-(p; p_1, p_2)$ are distributions affiliated with the “*generalized retarded four-point functions*” (see [8,9]).

Here again, any usable information on the support of the energy-momentum spectrum of the theory in the corresponding two-field sector will amount to specifying an open subset $\mathcal{R}$ in the space of energy-momentum vectors (p, p_1, p_2) whose *boundary only depend on the total energy-momentum vector* p in which the distributions $< \tilde{R}(p_1, p - p_1) \tilde{R}(p_2, -p - p_2) >$ and $< \tilde{R}(p_2, -p - p_2) \tilde{R}(p_1, p - p_1) >$ vanish together. In view of (4), such a support property (corresponding to the knowledge of the “intermediate states in the latter matrix elements”) then implies the *coincidence relation* $F|_{\mathcal{R}}^+ = F|_{\mathcal{R}}^-$.

Moreover, as in the case of propagators, the postulate of microcausality for the field $\Phi(x)$ implies *properties of analytic continuation* of the previous objects in *complex energy-momentum space*, which play a crucial role. These properties hold (together with polynomial bounds) in the tempered-distribution setting of Wightman-LSZ fields considered in [17]; however they could also be justified (with a corresponding release of the bounds) in the hyperfunction setting along the same lines as in Sec. 2.1. Even if the description of these properties is more complicated, due to the occurrence of *three complex energy-momenta* $k = p + iq$, $k_1 = p_1 + iq_1$, $k_2 = p_2 + iq_2$, the situation reproduces the case of propagators *as far as the total energy-momentum p is concerned*. In fact, F^+ and F^- are boundary values of holomorphic functions from tubes $\mathcal{T}^+, \mathcal{T}^-$ whose projections onto the space of *complex total energy-momentum* $k = p + iq$ are respectively $T^+ : q \in V^+$ and $T^- : q \in V^-$, so that formula (4) still appears (like (3)) as a discontinuity formula: it indicates that the discontinuity between the two holomorphic functions $F^+(k; k_1, k_2)$ and $F^-(k; k_1, k_2)$ is known to vanish on the set $\mathcal{R}$.

However we must describe more carefully the situation concerning the analyticity properties of these functions in the “*internal momenta*” k_1 and k_2 . First, it is clear from formula (5) that in view of the support property of the retarded product ($x' - x$ contained in $\overline{V}^+$), $\tilde{R}$ is the boundary value of an (operator-valued) analytic function $\tilde{R}(k', p - k')$ from the tube $k' = p' + iq' : q' \in V^+$ for all real p . Therefore the r.h.s. of Eq.(4) is the boundary value of a holomorphic function $\Delta F(p; k_1, k_2)$ of (k_1, k_2) in the tube Θ defined by the conditions $q_1 \in V^+, q_2 \in V^+$ for all real p .

Now it is also shown [8,9] that the domains of analyticity of F^+, F^- implied by microcausality are the tubes $\mathcal{T}^+$ and $\mathcal{T}^-$ defined by the following conditions:

$$\mathcal{T}^+ : q \in V^+, q_1 \in V^+, q_2 \in V^+ \quad (6)$$

$$\mathcal{T}^- : -q \in V^+, q + q_1 \in V^+, q + q_2 \in V^+ \quad (7)$$

and one easily checks that these two tubes admit precisely as *their common boundary* (at $q = 0$) the tube Θ for all real p . On the latter, there holds the following discontinuity formula for the boundary values of F^+ and F^- :

$$\Delta F(p; k_1, k_2) = F^+(p; k_1, k_2) - F^-(p; k_1, k_2). \quad (8)$$

The main geometrical difference with respect to the case of propagators is that the tubes $\mathcal{T}^+$ and $\mathcal{T}^-$ in the big complex (k, k_1, k_2) -space are *not opposite* as it is the case for T^+ and T^- in k -space. As a matter of fact, in view of (6) and (7), the union of the tubes $\mathcal{T}^+$ and $\mathcal{T}^-$ admits a convex hull $\tilde{\mathcal{T}}$ which is contained in the tube defined by the conditions $q_1 \in V^+$, $q_2 \in V^+$, $q + q_1 \in V^+$, $q + q_2 \in V^+$. Now in such a situation, and provided the *coincidence relation* $F|_{\mathcal{R}}^+ = F|_{\mathcal{R}}^-$ holds true, there exists a generalized version of the edge-of-the-wedge theorem [10], which states that $F^+(k; k_1, k_2)$ and $F^-(k; k_1, k_2)$ still admit a *common analytic continuation* $F(k; k_1, k_2)$. The latter is analytic in the union of $\mathcal{T}^+$, $\mathcal{T}^-$ and of a complex set $\mathcal{N}(\mathcal{R})$ of the following form: $\mathcal{N}(\mathcal{R})$ is the intersection of a complex neighborhood of $\mathcal{R}$ with the convex hull $\tilde{\mathcal{T}}$ of $\mathcal{T}^+ \cup \mathcal{T}^-$; in other words, F^+ and F^- “communicate analytically” through the complex set $\mathcal{N}(\mathcal{R})$ which is bordered by $\mathcal{R}$, although not being analytic anymore in $\mathcal{R}$ itself.

In the present situation, the open set $\mathcal{R}$ is always of the following “cylindric” form: p_1 and p_2 are arbitrary and p varies in an open set $\underline{\mathcal{R}}$ (namely the projection of $\mathcal{R}$ onto p -space). Then the equivalence of the following two statements (proved in [8,9]) deserves to be stressed:

- a) the boundary values of $F^+(k; k_1, k_2)$ and $F^-(k; k_1, k_2)$ coincide on $\mathcal{R}$,
- b) $\Delta F(p; k_1, k_2)$ vanishes as an analytic function of (k_1, k_2) in Θ for all p in $\underline{\mathcal{R}}$.

Property b) means that the “bridge” in which F^+ and F^- have a common analytic continuation contains not only the “small” set $\mathcal{N}(\mathcal{R})$ but the “large common face” defined by the conditions (k_1, k_2) in Θ for all p in $\underline{\mathcal{R}}$.

As in Sec. 2, one is then led to make use of an analytic completion procedure in order to enlarge the primitive (“non-natural”) set $\Sigma_{\mathcal{R}} = \mathcal{T}^+ \cup \mathcal{T}^- \cup \mathcal{N}(\mathcal{R})$, in which $F(k; k_1, k_2)$ is known to be analytic. It turns out that one can obtain results very similar to those of Sec 2, which reproduce the corresponding physical interpretations. In fact, the Properties A', B' and C' listed below can be seen as exact counterparts of the respective Properties A, B and C, since they involve identical regions (now called) $\underline{\mathcal{R}}$ and $\hat{\underline{\mathcal{R}}}$ in the space of the total energy-momentum p , while the additional analyticity properties with respect to the internal energy-momenta k_1 and k_2 are a remnant of microcausality in these variables.

i) Dispersion laws with subluminal velocities

Under the weak assumption that energy-positivity only holds in privileged Lorentz frames (see Sec 2-1 and ⁽⁴⁾), microcausality implies that all the hypersurfaces $\mathcal{M}_i$ and $\mathcal{M}$ representing respectively dispersion laws $p_0 = \omega_i(\vec{p})$ of one-particle states and the border of the continuous energy-momentum spectrum of “intermediate states in the matrix elements” $< \hat{R}(p_1, p - p_1) \hat{R}(p_2, -p - p_2) >$ have to be space-like hypersurfaces.

This follows from

Property A':

Let $\mathcal{R}_{a,b}$ be the set of all points (p, p_1, p_2) such that p belongs to a neighborhood of a given time-like segment $]a, b[$ with end-points $p = a$ and $p = b$ (b in the future of a). Then any function $F(k; k_1, k_2)$ holomorphic in $\Sigma_{\mathcal{R}_{a,b}} = \mathcal{T}^+ \cup \mathcal{T}^- \cup \mathcal{N}(\mathcal{R}_{a,b})$ admits an analytic continuation in a (complex) domain which contains the set of all points (p, k_1, k_2) such that p belongs to the double-cone $\diamond_a^b$ and (k_1, k_2) varies arbitrarily in the tube Θ .

The argument of Sec 2-1, based on the consideration of time-like segments $]a, b[$ with b contained in $\mathcal{M}$ or $\mathcal{M}_1$, then shows again the necessity of the space-like character of these hypersurfaces. In fact, for all such choices of $]a, b[$, the conclusion of Property A' implies that the discontinuity $\Delta F(p; k_1, k_2)$ of F vanishes for all p in $\diamond_a^b$ and (k_1, k_2) in Θ and therefore that the distribution $< [\tilde{R}(p_1, p - p_1), \tilde{R}(p_2, -p - p_2)]_{\pm} >$ vanishes for all p in $\diamond_a^b$ and (p_1, p_2) arbitrary.

ii) *Lorentz invariance of dispersion laws*

Under the (usual) strong assumption that energy-positivity holds in all Lorentz frames, microcausality implies (as in Sec 2-2) that all the hypersurfaces $\mathcal{M}_i$ and $\mathcal{M}$ representing respectively dispersion laws $p_0 = \omega_i(\vec{p})$ of one-particle states and the border of the continuous energy-momentum spectrum of "intermediate states in the matrix elements" $< \tilde{R}(p_1, p - p_1) \tilde{R}(p_2, -p - p_2) >$ have to be hyperboloid-shells with equations of the form $p_0 = \sqrt{\vec{p}^2 + m_i^2}$, $p_0 = \sqrt{\vec{p}^2 + M^2}$. This follows from the applicability of

Property B':

Let $\mathcal{R} = \mathcal{R}_0$ be the set of all (real) configurations (p, p_1, p_2) such that the total energy-momentum vector $p = (p_0, \vec{p})$ belongs to the following region $\underline{\mathcal{R}}_0 : |p_0| < |\vec{p}|$. Then any function $F(k; k_1, k_2)$ holomorphic in $\Sigma_{\mathcal{R}_0} = \mathcal{T}^+ \cup \mathcal{T}^- \cup \mathcal{R}_0$ admits an analytic continuation in the domain $\tilde{\Sigma}_{\mathcal{R}_0}$ which is the set of all complex configurations (k, k_1, k_2) belonging to the convex hull $\tilde{\mathcal{T}}$ of $\mathcal{T}^+ \cup \mathcal{T}^-$ and such that $k^2 \equiv k_0^2 - \vec{k}^2$ is different from any positive number and from zero,

supplemented by

Property C' (Lorentz-invariance of the borders of the spectral supports):

If $\mathcal{R}$ is any real open set, not necessarily connected, containing $\mathcal{R}_0$ and of "cylindric form" $p \in \underline{\mathcal{R}}$, with $\underline{\mathcal{R}} \supset \underline{\mathcal{R}}_0$, p_1, p_2 arbitrary, then every function $F(k; k_1, k_2)$ holomorphic in $\Sigma_{\mathcal{R}} = \mathcal{T}^+ \cup \mathcal{T}^- \cup \mathcal{R}$ admits an analytic continuation in the set of all configurations (p, k_1, k_2) such that (k_1, k_2) belongs to the tube Θ and p varies in an open set $\underline{\mathcal{R}}$ defined as in Property C: it is (for $d \geq 3$) the set of all real vectors p whose Minkowskian norm p^2 has a value already taken at some vector in $\underline{\mathcal{R}}$. Equivalently (but then including the case $d = 2$), it is the set of all vectors p obtained from vectors in $\underline{\mathcal{R}}$ by the action of a (real) Lorentz transformation.

The conclusion of Property C' implies that the discontinuity $\Delta F(p; k_1, k_2)$ of the holomorphic function $F(k; k_1, k_2)$ vanishes for all p in $\underline{\mathcal{R}}$ and (k_1, k_2) in Θ and therefore

that the distribution $\langle [\tilde{R}(p_1, p - p_1), \tilde{R}(p_2, -p - p_2)]_{\pm} \rangle$ vanishes for all p in $\hat{\mathcal{R}}$ and (p_1, p_2) arbitrary. It thus expresses the property of Lorentz invariance of the borders of the energy-momentum spectrum and therefore (according to the same analysis as in Sec 2-2) the results announced above follow.

A derivation of Properties A', B' and C' can be given along the same line as the proofs of Properties A, B and C presented above in Sec. 2. Let us only mention here that Property A' corresponds to a specific case of the double-cone theorem for tubes $\mathcal{T}^+$, $\mathcal{T}^-$ in general (i.e. non-opposite) situations (see [16]) and that Property B' is exactly the statement given in Theorem 1 of [8] for the case of $n = 3$ vector variables, with $m = 0$.

Remark: In the statements previously given under i) and ii), the constraints which were obtained concern the shape of the energy-momentum spectrum as it appears in the subspace of two-field states generated by retarded products of the following form $R[\varphi] \rangle = \int \varphi(x, x') \theta(x'_0 - x_0) [\Phi(x'), \Phi(x)]_{\pm} \rangle dx dx'$.⁽⁶⁾ However, it is clear that the same treatment and results are valid as well for two-field states generated by the corresponding advanced products, and therefore for the subspace generated by all states of the form $C[\varphi] \rangle = \int \varphi(x, x') [\Phi(x'), \Phi(x)]_{\pm} \rangle dx dx'$ (for all admissible test-functions φ).

The general case:

We shall now end this section by explaining why the previous treatment of spectral properties of the space of “two-field states” can be generalized to the spaces of “ n -field states” for all $n \geq 3$. Although it is not here the right place for presenting this general treatment with all its technical details, it is still possible to indicate briefly how it works.

The formalism of *generalized retarded operators* (g.r.o.) [17] allows one to introduce *generalized absorptive parts*: these are expectation values of (anti-) commutators of the following form $\langle [\tilde{R}_{\alpha}(\{p_i; i \in I\}), \tilde{R}_{\alpha'}(\{p'_{i'}; i' \in I'\})]_{\pm} \rangle$, where the operators $\tilde{R}_{\alpha}(\{p_i; i \in I\})$ and $\tilde{R}_{\alpha'}(\{p'_{i'}; i' \in I'\})$ denote the Fourier transforms of n -point g.r.o. R_{α} , $R_{\alpha'}$ with supports contained in relevant corresponding salient cones $\mathcal{C}_{\alpha}$ and $\mathcal{C}_{\alpha'}$ in the space of differences $x_j - x_k$ (resp. $x'_{j'} - x'_{k'}$) of space-time vectors: these cones are (non-trivial) analogs of the supports of the usual retarded and advanced operators of the case $n = 2$ (i.e. $x_1 - x_2 \in \bar{V}^{\pm}$). In our notation, I and I' represent disjoint subsets of n elements ($|I| = |I'| = n$) of the set $\{1, 2, \dots, 2n\}$ and the corresponding energy-momenta $p_i, p'_{i'}$ are linked by the energy-momentum conservation law $p = \sum_{i \in I} p_i = - \sum_{i' \in I'} p'_{i'}$, p being the total energy-momentum of the corresponding channel (I, I') of the $2n$ -point function of the fields considered; as previously (see ⁽³⁾), it is understood that the distribution $\delta(\{\sum_{i \in I} p_i\} + \{\sum_{i' \in I'} p'_{i'}\})$ has been factored out in the brackets $\langle \rangle$.

We then claim that for each n and each (I, I') there exists a complete set of g.r.o. R_{α} , $R_{\alpha'}$ whose Fourier transforms satisfy a discontinuity formula analogous to (4) of the

⁽⁶⁾ Rigorously speaking, the passage from support properties of the “scalar” distribution $\langle \tilde{R}(p_1, p - p_1) \tilde{R}(p_2, -p - p_2) \rangle$ to corresponding support properties of the vector-valued distribution $\varphi \rightarrow R[\varphi] \rangle$ relies on a Hilbert-space-norm argument.

following form

$$F_{\alpha,\alpha'}^+(\{p_i; i \in I\}; \{p'_{i'}; i' \in I'\}) - F_{\alpha,\alpha'}^-(\{p_i; i \in I\}; \{p'_{i'}; i' \in I'\}) = \\ < [\tilde{R}_\alpha(\{p_i; i \in I\}), \tilde{R}_{\alpha'}(\{p'_{i'}; i' \in I'\})]_{\pm} >; \quad (9)$$

in the latter, $F_{\alpha,\alpha'}^\pm(\{p_i; i \in I\}; \{p'_{i'}; i' \in I'\})$ are distributions affiliated with the “generalized retarded $2n$ -point functions” which are boundary values of analytic functions (still denoted by) $F_{\alpha,\alpha'}^\pm(\{k_i; i \in I\}; \{k'_{i'}; i' \in I'\})$ from respective tubes $\mathcal{T}_{\alpha,\alpha'}^+, \mathcal{T}_{\alpha,\alpha'}^-$, in the space of complex vectors $k_i = p_i + iq_i$, $k'_{i'} = p'_{i'} + iq'_{i'}$, such that $k = p + iq = \sum_{i \in I} k_i = -\sum_{i' \in I'} k'_{i'}$. These pairs of tubes play the same role as the pair $(\mathcal{T}^+, \mathcal{T}^-)$ of the case of two-field states: all points in $\mathcal{T}_{\alpha,\alpha'}^+$ (resp. $\mathcal{T}_{\alpha,\alpha'}^-$) satisfy the condition $q = \Im m k \in V^+$ (resp. V^-). Micro-causality is a basic ingredient in the proof of the previous statement, which relies on the results of [17 d), e)].

We are again led to express the energy-momentum spectral assumptions of the theory in the corresponding n -field sector by specifying an open subset $\mathcal{R}$ in the space of energy-momentum vectors $p_i, p'_{i'}$ whose boundary *only depend on the total energy-momentum vector* $p = \sum_{i \in I} p_i$, in which the distributions $< \tilde{R}_\alpha(\{p_i; i \in I\}) \tilde{R}_{\alpha'}(\{p'_{i'}; i' \in I'\}) >$ and $< \tilde{R}_{\alpha'}(\{p'_{i'}; i' \in I'\}) \tilde{R}_\alpha(\{p_i; i \in I\}) >$ vanish together. Here again, the edge-of-the-wedge theorem [10] implies that $F_{\alpha,\alpha'}^+$ and $F_{\alpha,\alpha'}^-$ have a common analytic continuation $F_{\alpha,\alpha'}$ in a set of the form $\Sigma_{\mathcal{R}} = \mathcal{T}_{\alpha,\alpha'}^+ \cup \mathcal{T}_{\alpha,\alpha'}^- \cup \mathcal{N}(\mathcal{R})$.

One could then present the “ n -field-state version” of Properties A', B' and C' in a way which closely parallels the two-field state case. For brevity, we shall not repeat the full statements and the corresponding physical interpretations which are identical to those listed above in paragraphs i) and ii) under the respective “weak” and “strong” forms of the energy-positivity condition. To exhibit the parallelism of the geometry of the n -field case with the one of the two-field case, it is sufficient to make a little more precise the description of the situation in the sets of energy-momentum vectors k_i and $k'_{i'}$, and the characterization of the domains $\mathcal{T}_{\alpha,\alpha'}^+, \mathcal{T}_{\alpha,\alpha'}^-$, and of their common face in the subspace $k = p$ real.

For p real, we introduce the sets of complex vectors $K_I = \{\underline{k}_i = k_i - \frac{p}{n}; i \in I\}$ and $K'_{I'} = \{\underline{k}'_{i'} = k'_{i'} + \frac{p}{n}; i' \in I'\}$ linked by the relations $\sum_{i \in I} \underline{k}_i = \sum_{i' \in I'} \underline{k}'_{i'} = 0$; correspondingly $Q_I = \Im m K_I$ (resp. $Q'_{I'} = \Im m K'_{I'}$) is the set of all q_i (resp. $q'_{i'}$) such that $\sum_{i \in I} q_i = 0$ (resp. $\sum_{i' \in I'} q'_{i'} = 0$). Each of the sets of vectors $K_I, K'_{I'}$ (resp. $Q_I, Q'_{I'}$) varies in a space of $(n-1)$ independent complex (resp. real) energy-momentum vectors.

By taking into account analogs of formula (5) for the operators $\tilde{R}_\alpha$ and $\tilde{R}_{\alpha'}$ together with linear identities between them (called “Steinmann relations” [17]), one can deduce from the support properties of R_α and $R_{\alpha'}$ (namely $\text{supp } R_\alpha \subset \mathcal{C}_\alpha$, $\text{supp } R_{\alpha'} \subset \mathcal{C}_{\alpha'}$) the following analyticity property: the r.h.s. of Eq.(9) is for every real p the boundary value of an analytic function $\Delta F_{\alpha,\alpha'}(p; K_I, K'_{I'})$ of $(K_I, K'_{I'})$, holomorphic in a well-defined tube $\Theta_{\alpha,\alpha'}$ (playing the same role as Θ in the case $n = 2$). This tube is specified by a set of conditions of the following type in the space of the imaginary parts $(Q_I, Q'_{I'})$. There exists a set Π_α of partitions (J, L) of I and a set $\Pi'_{\alpha'}$ of partitions (J', L') of I' such

that the defining conditions for $\Theta_{\alpha,\alpha'}$ are: $q_J = -q_L \in V^+$ and $q'_{J'} = -q'_{L'} \in V^+$ for all (J, L) in Π_α and all (J', L') in $\Pi'_{\alpha'}$: in the latter the notation q_J (resp. $q'_{J'}$) refers to the corresponding partial sum $\sum_{i \in J} q_i$ (resp. $\sum_{i' \in J'} q'_{i'}$). The sets Π_α and $\Pi'_{\alpha'}$ are not arbitrary but must satisfy the so-called “cell-conditions” (see [17]) which express the fact that no linear subspace with equation $q_M = 0$ or $q'_{M'} = 0$, with $M \subset I$ and $M' \subset I'$ intersects the domain $\Theta_{\alpha,\alpha'}$.

Now it can be shown that the tubes $\mathcal{T}_{\alpha,\alpha'}^+$ and $\mathcal{T}_{\alpha,\alpha'}^-$ in which the functions $F_{\alpha,\alpha'}^+$ and $F_{\alpha,\alpha'}^-$ are holomorphic are defined by the following conditions:

$$\mathcal{T}_{\alpha,\alpha'}^+ : \quad q \in V^+, \quad -q_L \in V^+ \text{ and } q'_{J'} \in V^+ \quad (10)$$

for all (J, L) in Π_α and all (J', L') in $\Pi'_{\alpha'}$;

$$\mathcal{T}_{\alpha,\alpha'}^- : \quad -q \in V^+, \quad q_J = -q_L + q \in V^+ \text{ and } q'_{J'} + q = -q'_{L'} \in V^+ \quad (11)$$

for all (J, L) in Π_α and all (J', L') in $\Pi'_{\alpha'}$.

These two tubes admit as their common boundary (at $q = 0$) the tube $\Theta_{\alpha,\alpha'}$ for all real p . On the latter, there holds the following discontinuity formula for the boundary values of $F_{\alpha,\alpha'}^+$ and $F_{\alpha,\alpha'}^-$:

$$\begin{aligned} \Delta F_{\alpha,\alpha'}(p; K_I, K'_{I'}) = \\ F_{\alpha,\alpha'}^+(\{k_i; i \in I\}; \{k'_{i'}; i' \in I'\})|_{q=0} - F_{\alpha,\alpha'}^-(\{k_i; i \in I\}; \{k'_{i'}; i' \in I'\})|_{q=0}. \end{aligned} \quad (12)$$

One easily checks that the defining conditions (10), (11) of the tubes $\mathcal{T}_{\alpha,\alpha'}^+$ and $\mathcal{T}_{\alpha,\alpha'}^-$ are completely analogous to the defining conditions (6), (7) of $\mathcal{T}^+$ and $\mathcal{T}^-$, up to the replacement of the two vector variables q_1, q_2 by all the vector variables $-q_L, q'_{J'}$ corresponding to the sets of partitions $\Pi_\alpha, \Pi_{\alpha'}$.

As a matter of fact, it is known (see [17]) that it is sufficient to consider a subset of g.r.o. called “Steinmann monomials” $R_\alpha, R_{\alpha'}$ for which each of the corresponding sets $\Pi_\alpha, \Pi_{\alpha'}$ contains *exactly* $n - 1$ partitions (one also says that the corresponding cell-conditions are “simplicial”); in fact, the most general g.r.o. are linear combinations of these Steinmann monomials. It then turns out that in this restricted class of g.r.o. the analog of Property B' coincides with Theorem 1 of [8] in its general n -vector form (with $m = 0$): this property states that *any function holomorphic in $\Sigma_{\mathcal{R}_0} = \mathcal{T}_{\alpha,\alpha'}^+ \cup \mathcal{T}_{\alpha,\alpha'}^- \cup \mathcal{N}(\mathcal{R}_0)$ (with $\mathcal{R}_0$ now defined by the conditions $|p_0| < |\vec{p}|$, K_I and $K'_{I'}$ real and arbitrary), admits an analytic continuation at all the points $(k, K_I, K'_{I'})$ in the convex hull of the tube $\mathcal{T}_{\alpha,\alpha'}^+ \cup \mathcal{T}_{\alpha,\alpha'}^-$ such that $k^2 \equiv k_0^2 - \vec{k}^2$ is different from any positive number and from zero. Property C' then follows from B' as in the case $n = 2$, while Property A' corresponds again to the double-cone theorem in a geometrical situation of general type.*

These considerations can be completed by a remark similar to the one given at the end of the case $n = 2$ (including footnote ⁽⁶⁾): since the g.r.o. generate (by linear combinations of Steinmann monomials) all the multiple (anti-)commutators of n field operators, the constraints on the energy-momentum spectrum apply to the subspace generated by all states

of the form $C[\varphi] \geq \int \varphi(x_1, \dots, x_{n-1}, x_n) [\Phi(x_1), [\dots, [\Phi(x_{n-1}, \Phi(x_n)) \dots]]] > dx_1 \dots dx_n$.
(for all admissible test-functions φ).

4 Concluding remarks

In this paper, we have displayed the geometrical constraints on the shape of the energy-momentum spectrum which result from microcausality together with (weak or strong) energy-positivity requirements in any (boson or fermion) interacting field theory. The Lorentz-invariant shape of the spectrum is therefore proven for possibly existing cases of field theories with Lorentz symmetry breaking with a high degree of generality. This is due to the purely geometrical character of our method, based on analyticity properties in several complex variables, which has allowed a strict exploitation of the latter requirements *in terms of the analytic Green's functions of the fields*: it is in terms of these objects that the spectral constraints are expressed and the degree of generality which is reached in this approach presents various aspects.

As already noticed (see our Remark in Sec. 3 and footnote ⁽⁶⁾), the Hilbert space interpretation of these constraints can be done separately. An advantage of the Green's function approach is therefore the fact that the constraints obtained are still proven to hold in an indefinite-metric framework, as for example in the usual treatment of the QCD-fields with a gauge-fixing preserving the microcausality conditions for the Green's functions.

Concerning the fact, already stressed above, that the results apply to the case of possible nonconventional field theories with arbitrarily wild short-distance singularities (i.e. of hyperfunction type, as specified in Sec. 2.1), we can say that the generality of the results goes even further: the latter are valid as soon as the Green's functions are analytic in the relevant tube domains of momentum space without any restriction on the increase of these functions at infinity, which can be considered as the most general expression of microcausality (note that mild exponential bounds in the energy were still obtained in the x -space hyperfunction setting, as noticed at the end of Sec. 2.1). In this connection, the method applies for example to the special case of Green's functions enjoying a very regular behaviour at large real energy-momenta, but exponentially increasing at purely imaginary energies: this is precisely the case for the Green's functions of the fields generated (via space-time translations) by local observables in the "local quantum physics" framework of [4], which were considered in the original works by Borchers and Buchholz [2,3] on the present subject.

References

- [1] V.A. Kostelecký and R. Lehnert, Phys. Rev. D **63**, 065008 (2001).
- [2] H.J. Borchers and D. Buchholz, Commun. Math. Phys. **97**, 169-185 (1985).
- [3] H.J. Borchers, Fizika (Zagreb) **17**, 289-304 (1985).
- [4] R. Haag, *Local Quantum Physics* (Springer-Verlag, Berlin-Heidelberg-New York, 1992).
- [5] H.J. Borchers, Commun. Math. Phys. **22**, 49-54 (1962).

- [6] H.J. Borchers, *Translation Group and Particle Representations in Quantum Field Theory* (Springer, New York, 1996).
- [7] R. Jost and H. Lehmann, *Nuovo Cimento* **5**, 1598-1610 (1957); F.J. Dyson, *Phys. Rev.* **110**, 1460-1464 (1958).
- [8] J. Bros , H. Epstein and V. Glaser, *Nuovo Cimento* **31** , 1265-1302 (1964).
- [9] H. Epstein, *Some analytic properties of scattering amplitudes in quantum field theory* in “Particle Symmetries and Axiomatic Field Theory”, Brandeis Summer School 1965,(Gordon and Breach, New York,1966).
- [10] H. Epstein, *Journ. Math. Phys.* **1**, 524-531 (1960).
- [11] S. Bochner and W.T. Martin, *Several complex variables* (Princeton University Press, Princeton, NJ, 1948).
- [12] A.S. Wightman, *Analytic functions of several complex variables* in “Relations de dispersion et particules élémentaires”, Les Houches Summer School 1960, C. De Witt and R. Omnes eds (Hermann, Paris, 1960).
- [13] L. Asgeirsson, *Math. Ann.* **113**, 321 (1936).
- [14] V.S. Vladimirov, *Doklady Akad. Nauk SSSR* **134**, 251 (1960).
- [15] H.J. Borchers, *Nuovo Cimento* **19**, 787-796 (1961).
- [16] J. Bros , H. Epstein, V. Glaser and R.Stora, in “Hyperfunctions and Theoretical Physics”, *Lecture Notes in Mathematics* **449**, 185 (Springer-Verlag, Berlin, 1975).
- [17] -a) D. Ruelle, *Nuovo Cimento* **19**, 356 (1961) and Thesis, Zurich 1959;
 -b) O. Steinmann, *Helv. Phys. Acta* **33**, 257 (1960); **33**, 347 (1960);
 -c) H.Araki and N. Burgoyne, *Nuovo Cimento* **18**, 342 (1960); H. Araki, *J. Math. Phys.* **2**, 163 (1961);
 -d) J. Bros, *Comptes-rendus RCP* no 25, CNRS Strasbourg (1967) and Thesis, Paris 1970;
 -e) H. Epstein, V. Glaser and R. Stora, *General Properties of the n-point Functions in Local Quantum Field Theory* in “Structural Analysis of Collision Amplitudes”, Les Houches June Institut 1975, R. Balian and D. Iagolnitzer eds (North-Holland, Amsterdam, 1976).
- [18] R. F. Streater and A. S. Wightman, *PCT, Spin and Statistics and all that*, Benjamin, New York (1964)
- [19] H. Lehmann, K. Symanzik and W. Zimmermann, *Nuovo Cimento* **1**, 205 (1955) and **6**, 319 (1957)
- [20] S. Nagamachi and N. Mugibayashi, *Letters in Math. Phys.* **1**, 259-264 (1976).
- [21] A. Jaffe, *Phys. Rev.* **158**, 1454 (1967).